

HOTLINE: Effective February 18, 2020

0055167 Kappa/Lambda Quantitative Free Light Chains with Ratio, Serum KAP/LAM F

Methodology: Quantitative Immunoturbidimetry

Performed: Sun-Sat **Reported:** 1-4 days

Specimen Required: Collect: Serum Separator Tube (SST).

Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer 2 mL serum to an ARUP

Standard Transport Tube. (Min: 0.5 mL) Storage/Transport Temperature: Refrigerated. Unacceptable Conditions: Plasma.

Stability (collection to initiation of testing): After separation from cells: Ambient: Unacceptable; Refrigerated: 3 weeks;

Frozen: 6 months

Reference Interval:

Components	Reference Interval
Lambda Quantitative Free Light Chains, Serum	Effective February 18, 2020 5.71-26.30 mg/L
Kappa Quantitative Free Light Chains, Serum	Effective February 18, 2020 3.30-19.40 mg/L
Kappa/Lambda Free Light Chain Ratio, Serum	0.26-1.65

CPT Code(s): 83520 x2

HOTLINE NOTE: There is a unit of measure change associated with this test.

Change the unit of measure for component 0055168, Kappa Qnt Free Light Chains from mg/dL to mg/L. Change the unit of measure for component 0055169, Lambda Qnt Free Light Chains from mg/dL to mg/L.