

HOTLINE: Effective May 20, 2019

New Test	3001561	Hypersensitivity Pneumonitis Extended Panel (Farmer's Lung Panel)	HYPEREXT
Click for Pricing	g		
Methodology: Performed:	Qualitative Immunodiffusion/Quantitative ImmunoCAP Fluorescent Enzyme Immunoassay Sun-Sat		
Reported:	3-7 days		
Specimen Required	l: Collect: Serum S	eparator Tube (SST).	
	Specimen Preparation: Separate from cells ASAP or within 2 hours of collection. Transfer two 2.5 mL aliquots of serum to individual ARUP Standard Transport Tubes. (Min: 1 mL Per aliquot) Storage/Transport Temperature: Refrigerated.		
	Unacceptable Co Stability (collection	<u>inditions:</u> Plasma. Contaminated, hemolyzed, or severely lipemic specimens. <u>ion to initiation of testing)</u> . After separation from cells: Ambient: 48 hours; Refrigerated: 2 v eated freeze/thaw cycles)	veeks; Frozen:
		• *	

Interpretive Data: Allergen results of 0.10-0.34 kU/L are intended for specialist use as the clinical relevance is undetermined. Even though increasing ranges are reflective of increasing concentrations of allergen-specific IgE, these concentrations may not correlate with the degree of clinical response or skin testing results when challenged with a specific allergen. The correlation of allergy laboratory results with clinical history and in vivo reactivity to specific allergens is essential. A negative test may not rule out clinical allergy or even anaphylaxis. See Compliance Statement A: www.aruplab.com/CS

Note: Testing includes antibodies directed at *Aspergillus funigatus #1, A. funigatus #2, A. funigatus #3, A. funigatus #6, A. flavus, Aureobasidium pullulans, Pigeon Serum, Micropolyspora faeni, Thermoactinomyces vulgaris #1, T. candidus,* and *Saccharomonospora viridis.*

CPT Code(s): 86003 x3; 86005; 86331 x6; 86606 x5

New York DOH Approved.

HOTLINE NOTE: Refer to the Test Mix Addendum for interface build information.