



<u>Procedure</u>	<u>Result</u>	<u>Units</u>	<u>Ref Interval</u>	<u>Accession</u>	<u>Collected</u>	<u>Received</u>	<u>Reported/</u> <u>Verified</u>
Smith/RNP (ENA) Ab, IgG	60 H	AU/mL	[0 - 40]	19-161-900083	10-Jun-19 11:05:00	10-Jun-19 11:05:00	10-Jun-19 11:12:38

10-Jun-19 11:05:00 Smith/RNP (ENA) Ab, IgG:
 INTERPRETIVE INFORMATION: Smith/RNP (ENA) Antibody, IgG

- 29 AU/mL or Less Negative
- 30 - 40 AU/mL Equivocal
- 41 AU/mL or Greater Positive

Smith/RNP antibodies are frequently seen in patients with mixed connective tissue disease (MCTD) and are also associated with other systemic autoimmune rheumatic diseases (SARDs) such as systemic lupus erythematosus (SLE), systemic sclerosis, and myositis. Antibodies targeting the Smith/RNP antigenic complex also recognize Smith antigens, therefore, the Smith antibody response must be considered when interpreting these results.

* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab