



<u>Procedure</u>	<u>Result</u>	<u>Units</u>	<u>Ref Interval</u>	<u>Accession</u>	<u>Collected</u>	<u>Received</u>	<u>Reported/</u> <u>Verified</u>
Blastomyces dermatitidis EIA, Interp	Detected *		[Not Detected]	19-164-900118	13-Jun-19 15:53:00	13-Jun-19 15:53:00	13-Jun-19 16:32:04
Blastomyces Antigen, Urine	50.00	ng/mL		19-164-900118	13-Jun-19 15:53:00	13-Jun-19 15:53:00	13-Jun-19 16:32:04

13-Jun-19 15:53:00 Blastomyces dermatitidis EIA, Interp:
 INTERPRETIVE INFORMATION: Blastomyces Ag Quant by ELISA,
 Urine

Less than 1.25 ng/ml = Not Detected
 1.25-200 ng/mL = Detected
 Greater than 200 ng/mL = Detected (above the limit of
 quantification)
 The quantitative range of this assay is 1.25-200 ng/mL. Antigen concentrations greater
 than 200 ng/mL fall outside the linear range of the assay and cannot be accurately
 quantified.

This EIA test should be used in conjunction with other diagnostic procedures, including
 microbiological culture, histological examination of biopsy samples, and/or radiographic
 evidence, to aid in the diagnosis of blastomycosis.

Cross-reactivity with other endemic mycoses (Histoplasma and Coccidioides) may occur.
 Positive test results should be correlated with other clinical findings and relevant
 exposure history

Test developed and characteristics determined by ARUP Laboratories. See Compliance
 Statement B: aruplab.com/CS

* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab