## ARUP LABORATORIES | aruplab.com

500 Chipeta Way, Salt Lake City, Utah 84108-1221

phone: 801-583-2787, toll free: 800-522-2787 Tracy I. George, MD, Chief Medical Officer Patient Report

Patient Age/Sex: 35 years Male

Testosterone, Free/Total	l by Received: 16-D	ec-21 15:27 Report/	Verified: 16-Dec-21 15:46
Procedure	Result	Units	Reference Interval
Testosterone by	1068 <sup>i1</sup>	ng/dL	300-1080
Immunoassay			
Sex Hormone Binding	78 H i2	nmol/L	17-56
Globulin			
Testosterone,Free	131 <sup>i3</sup>	pg/mL	47-244
Calculation			
Testosterone,	1.2 <sup>L</sup>	96	1.6-2.9
Percentage Free			

## Test Information

il: Testosterone by Immunoassay

INTERPRETIVE INFORMATION: Testosterone by Immunoassay

Testosterone immunoassays are both imprecise and inaccurate at low testosterone concentrations, such as those found in children and cisgender females. For these individuals, testing by mass spectrometry is recommended; refer to Testosterone (Adult Females, Children, or Individuals on Testosterone-Suppressing Hormone Therapy) (ARUP test code 0081058). Free or bioavailable testosterone measurements may provide supportive information.

For individuals on testosterone hormone therapy, refer to cisgender male reference intervals. No reference intervals have been established for males younger than 14 years or for cisgender females. For a complete set of all established reference intervals, refer to ltd.aruplab.com/Tests/Pub/0070130.

i2: Sex Hormone Binding Globulin

REFERENCE INTERVAL: Sex Hormone Binding Globulin

Access complete set of age- and/or gender-specific reference intervals for this test in the ARUP Laboratory Test Directory (aruplab.com).

i3: Testosterone, Free Calculation

INTERPRETIVE INFORMATION: Testosterone, Free Calculation

Free testosterone concentration is calculated using total testosterone (measured by immunoassay) and the binding constant of testosterone and sex hormone-binding globulin (SHBG). Testosterone immunoassays are both imprecise and inaccurate at low testosterone concentrations, such as those found in children and cisgender females. For these individuals, testing by mass spectrometry is recommended; refer to Testosterone, Free (Adult Females, Children, or Individuals on Testosterone-Suppressing Hormone Therapy) (ARUP test code 0081059).

For individuals on testosterone hormone therapy, refer to cisgender male reference intervals. No reference intervals have been established for males younger than 14

\*=Abnormal, #=Corrected, C=Critical, f=Result Footnote, H-High, i-Test Information, L-Low, t-Interpretive Text, @=Performing lab

Unless otherwise indicated, testing performed at:

**ARUP Laboratories** 

500 Chipeta Way, Salt Lake City, UT 84108

Laboratory Director: Tracy I. George, MD

**ARUP Accession:** 21-350-900173 **Report Request ID:** 15066510

**Printed:** 16-Dec-21 16:27

Page 1 of 2

## ARUP LABORATORIES | aruplab.com

500 Chipeta Way, Salt Lake City, Utah 84108-1221

phone: 801-583-2787, toll free: 800-522-2787 Tracy I. George, MD, Chief Medical Officer

Patient Report

Patient Age/Sex: 35 years Male

## Test Information

Testosterone, Free Calculation

years or for cisgender females. For a complete set of all established reference intervals, refer to ltd.aruplab.com/Tests/Pub/0070109.

\*=Abnormal, #=Corrected, C=Critical, f=Result Footnote, H-High, i-Test Information, L-Low, t-Interpretive Text, @=Performing lab

Unless otherwise indicated, testing performed at:

**ARUP Laboratories** 

500 Chipeta Way, Salt Lake City, UT 84108 Laboratory Director: Tracy I. George, MD

**ARUP Accession:** 

21-350-900173 Report Request ID: 15066510

Printed:

16-Dec-21 16:27

Page 2 of 2