Example Report

ARUP Laboratories
500 Chipeta Way – Salt Lake City, UT 84108

(800)522-2787 - www.aruplab.com Julio C. Delgado, M.D. M.S., Director of Laboratories Patient Age/Gender: Unknown Unknown Printed: 18-Mar-20 14:41:35

	_	_		Reported/
Procedure	Result	Units	Ref Interval	Accession Collected Received Verified
Parvovirus B19 Antibody IgM	1.98 H	IV	[<=0.90]	20-078-900139 18-Mar-20 18-Mar-20 18-Mar-20
				14:19:00 14:21:00 14:23:29

18-Mar-20 14:19:00 Parvovirus B19 Antibody IgM:
INTERPRETIVE INFORMATION: Parvovirus B19 Antibody, IgM
0.90 IV or less Negative - No significant
level of detectable Parvovirus
B19 IgM antibody.
0.91 - 1.09 IV Equivocal - Repeat testing in
7-21 days may be helpful.
1.10 IV or greater Positive - IgM antibody to
Parvovirus B19 detected which
may indicate a current or
recent infection. However, low
levels of IgM antibodies may
occasionally persist for more
than 12 months post-infection.

The best evidence for current infection is a significant change on two appropriately timed specimens, where both tests are done in the same laboratory at the same time.

Appearance of an IgM antibody response normally occurs 7 to 14 days after the onset of disease. Testing immediately post-exposure is of no value without a later convalescent specimen. A residual IgM response may be distinguished from early IgM response to infection by testing sera from patients three to four weeks later for changing levels of specific IgM antibodies.

* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab

Chart ID: 13648560 Page 1 of 1