



Procedure	Result	Units	Ref Interval	Accession	Collected	Received	Reported/ Verified
Anti-gp210 Antibody, IgG	25.0 H	Units	[0.0-24.9]	20-077-900082	17-Mar-20 12:15:00	17-Mar-20 12:15:00	17-Mar-20 12:32:07
Anti-spl00 Antibody, IgG	25.0 H	Units	[0.0-24.9]	20-077-900082	17-Mar-20 12:15:00	17-Mar-20 12:15:00	17-Mar-20 12:32:07

17-Mar-20 12:15:00 Anti-gp210 Antibody, IgG:
 REFERENCE INTERVAL: Anti-gp210 Antibody, IgG

- 20.0 Units or less.....Negative
- 20.1-24.9 Units.....Equivocal
- 25.0 Units or greater.....Positive

GP210 IgG antibodies can be detected in patients with primary biliary cholangitis (PBC) and may be of diagnostic relevance in a subset of patients with PBC who are negative for anti-mitochondrial antibodies (AMA). These antibodies have a relatively low sensitivity with excellent specificity for PBC. A negative result does not rule out PBC.

17-Mar-20 12:15:00 Anti-spl00 Antibody, IgG:
 REFERENCE INTERVAL: Anti-spl00 Antibody, IgG

- 20.0 Units or less.....Negative
- 20.1-24.9 Units.....Equivocal
- 25.0 Units or greater.....Positive

SP100 IgG antibodies can be detected in patients with primary biliary cholangitis (PBC) and may be of diagnostic relevance in a subset of patients with PBC who are negative for anti-mitochondrial antibodies (AMA). These antibodies have a relatively low sensitivity with excellent specificity for PBC. A negative result does not rule out PBC.

* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab