

Specimen Collected: 17-Mar-22 11:03

Circulating Immune Complex Panel		Received: 17-Mar-22 11:03	Report/Verified: 17-Mar-22 12:07
Procedure	Result	Units	Reference Interval
Circulating Immune Complex, C1q Binding	2.0 ⁱ¹	ug Eq/mL	0.0-3.9
Circulating Immune Complex, C3 fragments	16 ^{H i2}	ug Eq/mL	<=15

Test Information

i1: Circulating Immune Complex, C1q Binding
 INTERPRETIVE INFORMATION: Circulating Immune Complex, C1q Binding
 Less than or equal to 3.9 ug Eq/mL is considered negative for circulating complement binding immune complexes.

Circulating immune complexes may be found without any evident pathology and positive results do not necessarily implicate the immune complex in a disease process.

i2: Circulating Immune Complex, C3 fragments
 INTERPRETIVE INFORMATION: Circulating Immune Complex, C3 fragments
 Many autoimmune disorders, chronic infections and malignancies are associated with circulating immune complexes. Quantitation of immune complexes assists in staging immunologic disorders. Detection of circulating immune complexes is not essential to any specific diagnosis. Circulating immune complexes may be found without any evident pathology and positive results do not necessarily implicate immune complex-related disease process. Values between 15 and 20 ug Eq/mL are considered equivocal for Circulating Immune Complex, C3 fragments assay. Repeat- testing using a new specimen is recommended, if clinically indicated.

*=Abnormal, #=Corrected, C=Critical, f=Result Footnote, H-High, i-Test Information, L-Low, t-Interpretive Text, @=Performing lab

Unless otherwise indicated, testing performed at:

ARUP Laboratories

500 Chipeta Way, Salt Lake City, UT 84108

Laboratory Director: Tracy I. George, MD

ARUP Accession: 22-076-900061

Report Request ID: 15082614

Printed: 22-Mar-22 07:49

Page 1 of 1