ARUP LABORATORIES | aruplab.com

PATIENT REPORT

500 Chipeta Way, Salt Lake City, Utah 84108-1221

phone: 801-583-2787, toll free: 800-522-2787

Jonathan R. Genzen, MD, PhD, Chief Medical Officer

Patient Age/Sex: 37 years Female

Specimen Collected: 20-Mar-23 14:26

Drug Panel 9A, Screen Only, Urine	Received: 23-Mar-23	14:29	Report/Verified: 23-Mar-23 15:15
Procedure	Result	Units	Reference Interval
Creatinine, Urine	52.7	mg/dL	[20.0-400.0]
Alcohol, Urn, Screen	Negative	mg/dL	[Cutoff 40]
Amphetamines, Urn, Screen	Positive	ng/mL	[Cutoff 300]
Barbiturates, Urn, Screen	Negative	ng/mL	[Cutoff 200]
Benzodiazepines,Urn,Screen	Negative	ng/mL	[Cutoff 200]
Cocaine, Urn, Screen	Negative	ng/mL	[Cutoff 150]
MDMA, Urn, Screen	Negative	ng/mL	[Cutoff 500]
Methadone, Urn, Screen	Negative	ng/mL	[Cutoff 150]
Opiates, Urn, Screen	Negative	ng/mL	[Cutoff 300]
Oxycodone, Urn, Screen	Negative	ng/mL	[Cutoff 100]
Phencyclidine, Urn, Screen	Negative	ng/mL	[Cutoff 25]
Propoxyphene, Urn, Screen	Negative	ng/mL	[Cutoff 300]
THC, Urn, Screen	Positive	ng/mL	[Cutoff 50]
Drug Panel 9A, Screen Only -	See Note 11		
Comments			

Test Information

i1: Drug Panel 9A, Screen Only - Comments
INTERPRETIVE INFORMATION: Drug Panel 9A, Screen Only, Urine

This is a screening test only. False positive and false negative results can occur. Positive results are not automatically reflexed to a confirmatory test. Confirmation by GC/MS and/or LC-MS/MS must be requested separately. Confirmatory testing for drugs and/or drug classes detected by this screening test is recommended.

The absence of expected drug(s) and/or drug metabolite(s) may indicate non-compliance, inappropriate timing of specimen collection relative to drug administration, poor drug absorption, diluted/adulterated urine, or limitations of testing. The concentration at which the screening test can detect a drug or metabolite varies within a drug class. The concentration value must be greater than or equal to the cutoff to be reported as positive. Interpretive questions should be directed to the laboratory.

The following opioids are not detected in this test, but can be ordered separately: fentanyl, buprenorphine, meperidine, tramadol, and tapentadol. A comprehensive panel that includes these opioids is available or individual opioid testing can be ordered. Refer to aruplab.com for test information.

For medical purposes only; not valid for forensic use.

Unless otherwise indicated, testing performed at:

ARUP Accession:

Report Request ID

500 Chipeta Way, Salt Lake City, UT 84108

Laboratory Director: Jonathan R. Genzen, MD, PhD

 ARUP Accession:
 23-079-900242

 Report Request ID:
 17731669

 Printed:
 23-Mar-23 16:01

Page 1 of 1

^{*=}Abnormal, #=Corrected, C=Critical, f=Result Footnote, H-High, i-Test Information, L-Low, t-Interpretive Text, @=Performing lab