

Specimen Collected: 20-Mar-23 14:26

Drug Panel 9A, Urn, Scrn w/Rflx to Conf | Received: 23-Mar-23 14:29 | Report/Verified: 23-Mar-23 15:27

Procedure	Result	Units	Reference Interval
Creatinine, Urine	59.2	mg/dL	[20.0-400.0]
Alcohol, Urn, Screen	Negative	mg/dL	[Cutoff 40]
Amphetamines, Urn, Screen	Negative	ng/mL	[Cutoff 300]
Barbiturates, Urn, Screen	Negative	ng/mL	[Cutoff 200]
Benzodiazepines, Urn, Screen	Negative	ng/mL	[Cutoff 200]
Cocaine, Urn, Screen	Positive	ng/mL	[Cutoff 150]
Methadone, Urn, Screen	Negative	ng/mL	[Cutoff 150]
Opiates, Urn, Screen	Negative	ng/mL	[Cutoff 300]
Phencyclidine, Urn, Screen	Negative	ng/mL	[Cutoff 25]
Propoxyphene, Urn, Screen	Negative	ng/mL	[Cutoff 300]
THC, Urn, Screen	Negative	ng/mL	[Cutoff 50]
CDASU 9A Comments	See Note ⁱ¹		

Procedure	Result	Units	Reference Interval
Cocaine Met, Urine, Quant	>1000 ⁱ²	ng/mL	

Test Information

i1: CDASU 9A Comments

INTERPRETIVE INFORMATION: Drug Panel 9A, Urn, Scrn w/Rflx to Conf

The absence of expected drug(s) and/or drug metabolite(s) may indicate non-compliance, inappropriate timing of specimen collection relative to drug administration, poor drug absorption, diluted/adulterated urine, or limitations of testing. The concentration at which the screening test can detect a drug or metabolite varies within a drug class. Specimens for which drugs or drug classes are detected by the screen are reflexed to a second, more specific technology (GC/MS, GC/FID, and/or LC-MS/MS). The concentration value must be greater than or equal to the cutoff to be reported as positive. Interpretive questions should be directed to the laboratory.

For medical purposes only; not valid for forensic use.

Oxycodone results are reported with the opiates results. MDMA results are reported with the amphetamines results. The following opioids are not detected in this test: fentanyl, buprenorphine, meperidine, tramadol, and tapentadol. A comprehensive panel that includes these opioids is available or individual opioid testing can be ordered. Refer to aruplab.com for test information.

i2: Benzoyllecgonine, Urn, Quant

INTERPRETIVE INFORMATION: Cocaine Metabolite, Urine,
Quantitative

*=Abnormal, #=Corrected, C=Critical, f=Result Footnote, H-High, i-Test Information, L-Low, t-Interpretive Text, @=Performing lab

Unless otherwise indicated, testing performed at:

ARUP Laboratories

500 Chipeta Way, Salt Lake City, UT 84108

Laboratory Director: Jonathan R. Genzen, MD, PhD

ARUP Accession: 23-079-900245

Report Request ID: 17731684

Printed: 23-Mar-23 16:13

Page 1 of 2

Test Information

i2: Benzoylcegonine, Urn, Quant

Methodology: Quantitative Gas Chromatography-Mass Spectrometry/Quantitative Liquid Chromatography-Tandem Mass Spectrometry.

Positive cutoff: 50 ng/mL

For medical purposes only; not valid for forensic use.

The concentration value must be greater than or equal to the cutoff to be reported as positive. Interpretive questions should be directed to the laboratory.

This test was developed and its performance characteristics determined by ARUP Laboratories. It has not been cleared or approved by the US Food and Drug Administration. This test was performed in a CLIA certified laboratory and is intended for clinical purposes.

*=Abnormal, #=Corrected, C=Critical, f=Result Footnote, H=High, i=Test Information, L=Low, t=Interpretive Text, @=Performing lab

Unless otherwise indicated, testing performed at:

ARUP Laboratories

500 Chipeta Way, Salt Lake City, UT 84108

Laboratory Director: Jonathan R. Genzen, MD, PhD

ARUP Accession: 23-079-900245

Report Request ID: 17731684

Printed: 23-Mar-23 16:13

Page 2 of 2