



<u>Procedure</u>	<u>Result</u>	<u>Units</u>	<u>Ref Interval</u>	<u>Accession</u>	<u>Collected</u>	<u>Received</u>	<u>Reported/</u> <u>Verified</u>
Hours Collected	24	hr		17-255-900027	12-Sep-17	12-Sep-17	12-Sep-17
Total Volume	1890	mL		17-255-900027	07:02:00	07:02:00	07:05:57
Creatinine, Urine - per volume	45	mg/dL		17-255-900027	12-Sep-17	12-Sep-17	12-Sep-17
Creatinine, Urine - per 24h	850	mg/d	[500-1,400]	17-255-900027	07:02:00	07:02:00	07:05:57
Copper, Urine - per volume	<b>3.3 H</b>	ug/dL	[0.3-3.2]	17-255-900027	12-Sep-17	12-Sep-17	12-Sep-17
Copper, Urine - per 24h	<b>62.4 H</b>	ug/d	[3.0-45.0]	17-255-900027	07:02:00	07:02:00	07:05:57
Copper, Urine - ratio to CRT	<b>73.3 H</b>	ug/g CRT	[10.0-45.0]	17-255-900027	12-Sep-17	12-Sep-17	12-Sep-17
					07:02:00	07:02:00	07:05:57

12-Sep-17 07:02:00 Copper, Urine - per volume:  
 INTERPRETIVE INFORMATION: Copper, Urine

Individuals with symptomatic Wilson disease usually excrete more than 100 ug copper per day. Other conditions associated with elevated urine copper include cholestatic liver disease, proteinuria, some medications, and contaminated specimens.

Although random specimens may contain diagnostic information, a 24-hour collection is a more consistent indicator of urine copper.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS

\* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab