



<u>Procedure</u>	<u>Result</u>	<u>Units</u>	<u>Ref Interval</u>	<u>Accession</u>	<u>Collected</u>	<u>Received</u>	<u>Reported/</u> <u>Verified</u>
Creatinine, Urine - per volume	250	mg/dL		17-255-900045	12-Sep-17 08:13:00	12-Sep-17 08:13:00	12-Sep-17 08:14:59
Cadmium, Urine - per volume	8.2 H	ug/L	[0.0-1.0]	17-255-900045	12-Sep-17 08:13:00	12-Sep-17 08:13:00	12-Sep-17 08:14:59
Cadmium Rnd Urn ratio/CRT nonoccupation	3.3 H	ug/g CRT	[0.0-3.2]	17-255-900045	12-Sep-17 08:13:00	12-Sep-17 08:13:00	12-Sep-17 08:14:59

12-Sep-17 08:13:00 Cadmium, Urine - per volume:
 INTERPRETATION INFORMATION: Cadmium, Urine

Urine cadmium levels can be used to assess cadmium body burden. In chronic exposures, the kidneys are the primary target organ. Symptoms associated with cadmium toxicity vary based upon route of exposure and may include tubular proteinuria, fever, headache, dyspnea, chest pain, conjunctivitis, rhinitis, sore throat and cough. Ingestion of cadmium in high concentration may cause vomiting, diarrhea, salivation, cramps, and abdominal pain.

See Compliance Statement B: aruplab.com/CS

* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab