



<u>Procedure</u>	<u>Result</u>	<u>Units</u>	<u>Ref Interval</u>	<u>Accession</u>	<u>Collected</u>	<u>Received</u>	<u>Reported/</u> <u>Verified</u>
Creatinine, Urine - per volume	55	mg/dL		17-257-900214	14-Sep-17 13:35:00	14-Sep-17 13:35:00	14-Sep-17 13:36:11
Copper, Urine - per volume	<1.0	ug/dL	[0.3-3.2]	17-257-900214	14-Sep-17 13:35:00	14-Sep-17 13:35:00	14-Sep-17 13:36:11
Copper, Urine - ratio to CRT	<18.2	ug/g CRT	[10.0-45.0]	17-257-900214	14-Sep-17 13:35:00	14-Sep-17 13:35:00	14-Sep-17 13:36:11

14-Sep-17 13:35:00 Copper, Urine - per volume:
 INTERPRETIVE INFORMATION: Copper, Urine

Individuals with symptomatic Wilson disease usually excrete more than 100 ug copper per day. Other conditions associated with elevated urine copper include cholestatic liver disease, proteinuria, some medications, and contaminated specimens.

Although random specimens may contain diagnostic information, a 24-hour collection is a more consistent indicator of urine copper.

Test developed and characteristics determined by ARUP Laboratories. See Compliance Statement B: aruplab.com/CS

* Abnormal, # = Corrected, C = Critical, f = Footnote, H = High, L = Low, t = Interpretive Text, @ = Reference Lab