Procedure HIV-1 Qnt by NAAT (copies/mL)	Result 47	<u>Units</u> <u>Ref Interva</u> cpy/mL	Reported/ <u>Accession</u> <u>Collected Received</u> <u>Verified</u> 18-244-900030 01-Sep-18 01-Sep-18 14-Sep-18 09:47:00 09:47:00 17:55:19
HIV-1 Qnt by NAAT (log copies/mL)	1.68 f	log cpy/mL	18-244-900030 01-Sep-18 01-Sep-18 14-Sep-18 09:47:00 09:47:00 17:55:19
HIV-1 Qnt by NAAT Interp	Detected *	[Not Detected]	18-244-900030 01-Sep-18 01-Sep-18 14-Sep-18 09:47:00 09:47:00 17:55:19

01-Sep-18 09:47:00 HIV-1 Qnt by NAAT (log copies/mL):

Human Immunodeficiency Virus 1 by Quantitative NAAT result was less than 500 copies/mL; therefore no further testing added.

01-Sep-18 09:47:00 HIV-1 Qnt by NAAT Interp:

INTERPRETIVE INFORMATION: HIV-1 by Quantitative NAAT, Plasma

Normal range for this assay is "Not Detected". The quantitative range of this assay is 1.47-7.00 log copies/mL (30-10,000,000 copies/mL).

An interpretation of "Not Detected" does not rule out the presence of inhibitors or HIV-1 RNA concentration below the level of detection of the assay. Care should be taken in the interpretation of any single viral load determination. The clinical significance of changes in HIV-1 RNA concentration has not been fully established; however, a change of 0.5 log copies/mL may be significant.

This assay should not be used for blood donor screening, associated re-entry protocols, or for screening Human Cell, Tissues and Cellular Tissue-Based Products (HCT/P).