

**Specimen Collected: 07-Sep-21 17:08**

Cardiolipin Antibody, IgG		Received: 07-Sep-21 17:09	Report/Verified: 07-Sep-21 17:15
Procedure	Result	Units	Reference Interval
Cardiolipin Antibody IgG	20 <sup>H</sup> <sup>i1</sup>	GPL	<=14

**Test Information**

i1: Cardiolipin Antibody IgG

INTERPRETIVE INFORMATION: Anti-Cardiolipin IgG Ab

- 0-14 GPL: Negative
- 15-19 GPL: Indeterminate
- 20-80 GPL: Low to Moderately Positive
- 81 GPL or above: High Positive

The persistent presence of IgG and/or IgM cardiolipin (CL) antibodies in moderate or high levels (greater than 40 GPL and/or greater than 40 MPL units or greater than 99th percentile) is a laboratory criterion for the diagnosis of antiphospholipid syndrome (APS). Persistence is defined as moderate or high levels of IgG and/or IgM CL antibodies detected in two or more specimens drawn at least 12 weeks apart (J Throm Haemost. 2006;4:295-306). Lower positive levels of IgG and/or IgM CL antibodies (above cutoff but less than 40 GPL and/or less than 40 MPL units) may occur in patients with the clinical symptoms of APS; therefore, the actual significance of these levels is undefined. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

\*=Abnormal, #=Corrected, C=Critical, f=Result Footnote, H-High, i-Test Information, L-Low, t-Interpretive Text, @=Performing lab

**Unless otherwise indicated, testing performed at:**

**ARUP Laboratories**

500 Chipeta Way, Salt Lake City, UT 84108

Laboratory Director: Tracy I. George, MD

**ARUP Accession:** 21-250-900135

**Report Request ID:** 15046422

**Printed:** 08-Sep-21 09:41