

Specimen Collected: 07-Sep-21 17:08

Cardiolipin Antibody, IgM		Received: 07-Sep-21 17:09	Report/Verified: 07-Sep-21 17:15
Procedure	Result	Units	Reference Interval
Cardiolipin Antibody IgM	20 ^H ⁱ¹	MPL	<=12

Test Information

i1: Cardiolipin Antibody IgM
 INTERPRETIVE INFORMATION: Anti-Cardiolipin IgM

0-12 MPL: Negative
 13-19 MPL: Indeterminate
 20-80 MPL: Low to Moderately Positive
 81 MPL or above: High Positive

The persistent presence of IgG and/or IgM cardiolipin (CL) antibodies in moderate or high levels (greater than 40 GPL and/or greater than 40 MPL units or greater than 99th percentile) is a laboratory criterion for the diagnosis of antiphospholipid syndrome (APS). Persistence is defined as moderate or high levels of IgG and/or IgM CL antibodies detected in two or more specimens drawn at least 12 weeks apart (J Throm Haemost. 2006;4:295-306). Lower positive levels of IgG and/or IgM CL antibodies (above cutoff but less than 40 GPL and/or less than 40 MPL units) may occur in patients with the clinical symptoms of APS; therefore, the actual significance of these levels is undefined. Results should not be used alone for diagnosis and must be interpreted in light of APS-specific clinical manifestations and/or other criteria phospholipid antibody tests.

*=Abnormal, #=Corrected, C=Critical, f=Result Footnote, H=High, i=Test Information, L=Low, t=Interpretive Text, @=Performing lab

Unless otherwise indicated, testing performed at:

ARUP Laboratories

500 Chipeta Way, Salt Lake City, UT 84108

Laboratory Director: Tracy I. George, MD

ARUP Accession: 21-250-900134**Report Request ID:** 15046420**Printed:** 08-Sep-21 09:41

Page 1 of 1