

Noninvasive Prenatal Aneuploidy Screening Test



PATIENTS.ANSWERS.RESULTS.®

For a complete list of
our **medical directors**,
please visit:



For a complete and
updated list of our **tests**,
please visit:



To learn more about
our **genetic counselors**,
please visit:



3003043 Non-Invasive Prenatal Aneuploidy Screen by Cell-Free DNA Sequencing

Prenatal cell-free DNA (cfDNA) screening, previously referred to as noninvasive prenatal testing (NIPT) or noninvasive prenatal screening (NIPS), tests for specific chromosomal aneuploidies in the fetus and may be performed as early as 10 weeks gestational age through the third trimester. It is the most sensitive and specific aneuploidy screening test. This is not a genetic diagnostic test.

The American College of Obstetricians and Gynecologists (ACOG), the American College of Medical Genetics and Genomics (ACMG), and the Society for Maternal-Fetal Medicine (SMFM) recommend that healthcare providers discuss prenatal genetic screening and diagnostic options with all pregnant patients, regardless of the pregnant person's age or fetal risk of chromosomal abnormality.^{1,2,3} ARUP offers both prenatal cfDNA screening and maternal serum screening, as well as comprehensive prenatal diagnostic options.

Conditions and characteristics screened by prenatal cfDNA screening:

- Trisomy 21 (Down syndrome)
- Trisomy 18 (Edwards syndrome)
- Trisomy 13 (Patau syndrome)
- Monosomy X (Turner syndrome)
- XXY (Klinefelter syndrome)
- XXX (Triple X syndrome)
- XYY
- Fetal sex (patient may opt out)

ARUP offers genetic counselor support to providers to help them understand prenatal cfDNA screening and its applications. The [Prenatal Testing for Chromosomal Abnormalities and Neural Tube Defects](#) topic on ARUP Consult[®] provides detailed information about recent clinical practice guidelines, the advantages and disadvantages of testing choices, and recommendations for discussing these complex options with patients.

ARUP's prenatal cfDNA screening reports emphasize the screening nature of this test and are designed with a focus on clarity and brevity. Reports provide a clear summary of the high- or low-risk screening result, with recommendations for follow-up. Important details such as positive predictive value, negative likelihood ratio, and relevant data from multicenter studies may be included. Fetal fraction is listed unless testing does not yield a result. In cases without results, an explanation of possible cause(s) and recommendations are included.

Which platform does ARUP's prenatal cfDNA screen utilize?

ARUP's prenatal cfDNA screen utilizes a massively parallel whole genome sequencing method developed by Illumina (Verinata).⁴ The likelihood of aneuploidy is based on an over- or underrepresentation of interrogated chromosomes.



Prenatal Testing
for Chromosomal
Abnormalities and
Neural
Tube Defects



Performance of prenatal cfDNA screening

In 2017, a well-respected meta-analysis of prenatal cfDNA screening performance found that “there was no obvious difference in performance” between the four methodologies used for cfDNA prenatal aneuploidy screening.⁵ Multiple large studies have demonstrated that the Illumina prenatal cfDNA screening platform is robust, and ARUP has validated every aspect of the testing process.^{6,7,8} The analytic validation accuracy study results demonstrate that ARUP's prenatal cfDNA screening analysis and interpretation are consistent with the accuracy reported in large data sets published based on clinical studies utilizing this platform. Detailed information about test performance can be found in the [Noninvasive Prenatal Aneuploidy Screen by Cell-Free DNA Sequencing Test Fact Sheet](#).

Why choose ARUP for prenatal cfDNA screening?

- Dependable quality from a well-established, nonprofit, university-affiliated national reference laboratory
- Support provided to clinicians and clients by an integrated team of board-certified laboratory geneticists and genetic counselors who specialize in prenatal genetics and who can aid providers in test selection and results interpretation
- Opportunity for integration with a comprehensive test menu that includes a suite of testing related to women's health (e.g., carrier screening, maternal serum screening, prenatal cfDNA screening, and prenatal diagnostic testing)
- Succinct, clear reports that include recommendations and data to support providers
- Integration with the healthcare system
 - Results that interface with EMR and have links to enhanced reports
 - Straightforward billing consistent with nongenetic lab tests
 - Standard processing and sendout of samples, consistent with other sendout lab orders

Published Clinical Sensitivity and Specificity^{a,b}

Condition	Clinical Sensitivity	Clinical Specificity	PPV ^c	NPV
Trisomy 21	98.9%	99.7%	20.9-95.1%	>99.9%
Trisomy 18	>99.9%	99.8%	6.2-94.1%	>99.9%
Trisomy 13	>99.9%	99.8%	1.6-76.8%	>99.9%
Monosomy X	>99.9%	>99.8%	—	—

^a Male/female concordance is >99%.

^b Data are not sufficient to calculate sensitivity, specificity, PPV, and NPV for every condition.

^c PPV ranges are calculated based on sensitivity and specificity in Borth et al (2021)⁸ and the prevalence in a low-risk group (20 years of age, 30 weeks gestational age) through the prevalence in a high-risk group (44 years of age, 10 weeks gestational age).^{9,10} PPV is greatly affected by an individual's pretest risk for each of the screened conditions, which varies based on gestational age and maternal age.

NPV, negative predictive value; PPV, positive predictive value

Detailed information about PPV at increased pretest risk levels, various gestational ages, and maternal ages can be found in the [supplementary tables](#).

Healthcare providers with questions may contact an ARUP genetic counselor at 800-242-2787 ext. 2141.

For more information about cfDNA, visit: aruplab.com/cfDNAScreen

References

1. Screening for fetal chromosomal abnormalities: ACOG Practice Bulletin Summary, Number 226. *Obstet Gynecol.* 2020;136(4):859-867. <https://pubmed.ncbi.nlm.nih.gov/32976375/>
2. Gregg AR, Skotko BG, Benkendorf JL, et al. Noninvasive prenatal screening for fetal aneuploidy, 2016 update: a position statement of the American College of Medical Genetics and Genomics. *Genet Med.* 2016;18(10):1056-1065. <https://pubmed.ncbi.nlm.nih.gov/27467454/>
3. Driscoll DA, Gross SJ; Professional Practice and Guidelines Committee. First trimester diagnosis and screening for fetal aneuploidy. *Genet Med.* 2008;10(1):73-75. <https://pubmed.ncbi.nlm.nih.gov/18197059/>
4. Illumina. NIPT: Elevating the standard for prenatal screening. Accessed Jan 2026. <https://www.illumina.com/clinical/reproductive-genetic-health/nipt.html>.
5. Gil MM, Accurti V, Santacruz B, et al. Analysis of cell-free DNA in maternal blood in screening for aneuploidies: updated meta-analysis. *Ultrasound Obstet Gynecol.* 2017;50(3):302-314. <https://pubmed.ncbi.nlm.nih.gov/28397325/>
6. Cirigliano V, Ordoñez E, Rueda L, et al. Performance of the neoBona test: a new paired-end massively parallel shotgun sequencing approach for cell-free DNA-based aneuploidy screening. *Ultrasound Obstet Gynecol.* 2017;49(4):460-464. <https://pubmed.ncbi.nlm.nih.gov/27981672/>
7. Pertile MD, Flowers N, Vavrek D, et al. Performance of a paired-end sequencing-based noninvasive prenatal screening test in the detection of genome-wide fetal chromosomal anomalies. *Clin Chem.* 2021;67(9):1210-1219. <https://pubmed.ncbi.nlm.nih.gov/34077512/>
8. Borth H, Teubert A, Glaubitz R, et al. Analysis of cell-free DNA in a consecutive series of 13,607 routine cases for the detection of fetal chromosomal aneuploidies in a single center in Germany. *Arch Gynecol Obstet.* 2021;303(6):1407-1414. <https://pubmed.ncbi.nlm.nih.gov/33151425/>
9. Snijders RJ, Sundberg K, Holzgreve W, et al. Maternal age- and gestation-specific risk for trisomy 21. *Ultrasound Obstet Gynecol.* 1999;13(3):167-170. <https://pubmed.ncbi.nlm.nih.gov/10204206/>
10. Snijders RJ, Sebire NJ, Nicolaides KH. Maternal age and gestational age-specific risk for chromosomal defects. *Fetal Diagn Ther.* 1995;10(6):356-367. <https://pubmed.ncbi.nlm.nih.gov/8579773/>



A nonprofit enterprise of the University of Utah and its Department of Pathology

ARUP LABORATORIES

500 Chipeta Way
Salt Lake City, UT 84108-1221
Phone: 800-522-2787
Fax: 801-583-2712
aruplab.com | arupconsult.com

© 2026 ARUP Laboratories
MKT-TSTMRK-PP-026, Rev 00, January 2026

